

Contents

Contributors

xxii

Acknowledgements

xxiv

PART 1 Neurons: Excitable and Secretory Cells that Establish Synapses

Chapter 1 *Neurons*

C. Hammond

1.1	Neurons have a cell body from which emerge two types of processes: the dendrites and the axon	3
1.1.1	The somatodendritic tree is the neuron's receptive pole	3
1.1.2	The axon and its collaterals are the neuron's transmitter pole	5
1.2	Neurons are highly polarized cells with a differential distribution of organelles and proteins	7
1.2.1	The soma is the main site of macromolecule synthesis	7
1.2.2	The dendrites contain free ribosomes and synthesize some of their proteins.	9
1.2.3	The axon, to a large extent, lacks the machinery for protein synthesis	11
1.3	Axonal transport allows bidirectional communication between the cell body and the axon terminals	11
1.3.1	Demonstration of axonal transport	11
1.3.2	Fast anterograde axonal transport is responsible for the movement of membranous organelles from cell body towards axon terminals, and allows renewal of axonal proteins	12
1.3.3	Retrograde axonal transport is responsible for the movement of membranous organelles back from axon terminals to the cell body	15
1.3.4	Slow anterograde axonal transport moves cytoskeletal proteins and cytosoluble proteins	16
1.3.5	Axonal transport of mitochondria allows the turnover of mitochondria in axons and axon terminals	17
1.4	Neurons connected by synapses form networks or circuits	17
1.4.1	The circuit of the withdrawal medullary reflex	17
1.4.2	The spinothalamic tract or anterolateral pathway is a somatosensory pathway	18
1.5	Summary: the neuron is an excitable and secretory cell presenting an extreme functional regionalization	18
Appendix 1.1	The cytoskeletal elements in neurons	21
Further reading		22

Chapter 2 *Neuron–Glial Cell Cooperation*
C. Hammond

2.1	Astrocytes form a vast cellular network or syncytium between neurons, blood vessels and the surface of the brain	24
2.1.1	Astrocytes are star-shaped cells characterized by the presence of glial filaments in their cytoplasm	24
2.1.2	Astrocytes maintain the blood brain barrier in the adult brain	25
2.1.3	Astrocytes regulate the ionic composition of the extracellular fluid	26
2.1.4	Astrocytes take part in the neurotransmitter cycle	26
2.2	Oligodendrocytes form the myelin sheaths of axons in the central nervous system and allow the clustering of Na⁺ channels at nodes of Ranvier	26
2.2.1	Processes of interfascicular oligodendrocytes electrically isolate segments of central axons by forming the lipid-rich myelin sheath	27
2.2.2	Myelination enables rapid conduction of action potentials for two reasons	29
2.3	Microglia: ramified microglial cells represent the quiescent form of microglial cells in the central nervous system; they transform upon injury	30
2.3.1	Ramified microglial cells have long meandering processes	30
2.3.2	Do adult microglial cells play a role in immune processes?	31
2.4	Ependymal cells constitute an active barrier between blood and cerebrospinal fluid	31
2.4.1	Ependymal cells form an epithelium at the surface of the ventricles	31
2.4.2	Ependymal cells of the choroid plexus	31
2.4.3	Extrachoroidal ependymal cells	32
2.5	Schwann cells are the glial cells of the peripheral nervous system; they form the myelin sheath of axons or encapsulate neurons	33
2.5.1	Myelinating Schwann cells make the myelin sheath of peripheral axons	33
2.5.2	Non-myelinating Schwann cells encapsulate the axons and cell bodies of peripheral neurons	33
	Further reading	33

Chapter 3 *Ionic Fluxes Across the Neuronal Plasma Membrane*
C. Hammond

3.1	Observation and questions	36
3.1.1	There is an unequal distribution of ions across neuronal plasma membrane	36
3.1.2	Neuronal plasma membrane is permeable to ions, allowing both passive and active transport of ions	37
3.1.3	There is a difference of potential between the two faces of the membrane, called membrane potential	37
3.1.4	Questions	38
3.2	Na⁺, K⁺, Ca²⁺ and Cl⁻ ions passively cross the plasma membrane through transmembrane proteins – the channels	39
3.2.1	Channels are a particular class of transmembrane proteins	39
3.2.2	Voltage-gated channels open in response to a change in membrane potential	40

3.2.3	Ligand-gated channels opened by extracellular ligands, and receptor channels opened by neurotransmitters	41
3.2.4	Ligand-gated channels opened by intracellular ligands	41
3.2.5	Mechanically gated channels opened by mechanical stimuli	43
3.2.6	Other channels: junctional channels or gap junctions	43
3.2.7	Distribution of the various channels in the neuronal plasma membrane	44
3.3	The diffusion of ions through an open channel: What is an electrochemical gradient and an ionic current?	47
3.3.1	The structure of the channel pore determines the type of ion(s) that diffuse passively through the channel	47
3.3.2	The electrochemical gradient for a particular ion determines the direction of the passive diffusion of this ion through an open channel	48
3.3.3	The passive diffusion of ions through an open channel is a current	50
3.4	Active transport of Na^+, K^+, Ca^{2+} and Cl^- ions by pumps and transporters maintain the unequal distribution of ions	52
3.4.1	Pumps are ATPases that actively transport ions	52
3.4.2	Transporters use the energy stored in the transmembrane electrochemical gradient of Na^+ or other ions H^+	52
3.5	Summary	53
Appendix 3.1	Hydrophobicity profile of a transmembrane protein	54
Appendix 3.2	The Nernst equation	55
	Further reading	56

Chapter 4 *Basic Properties of Excitable Cells at Rest*

A. Nistri and A. Gutman

4.1	Ionic channels open at rest determine the resting membrane potential	57
4.1.1	The plasma membrane separates two media of different ionic composition	58
4.1.2	At rest most of the channels open are K^+ channels	58
4.1.3	In muscle cells, K^+ and Cl^- ion movements participate equally in resting membrane potential	59
4.1.4	In central neurons, K^+ , Cl^- and Na^+ ion movements participate in resting membrane potential: the Goldman–Hodgkin–Katz equation	60
4.1.5	Some principles related to the derivation of the GHK equation	61
4.2	Membrane pumps are responsible for keeping constant the concentration gradients across membranes	61
4.3	A simple equivalent electrical circuit for resting membrane properties	62
4.3.1	Membrane potential has an ohmic behaviour at rest	62
4.3.2	Stability, bistability and instability of resting membrane potential	64
4.3.3	An electrical model of resting membrane potential	64
4.4	Advantages and disadvantages of sharp (intracellular) versus patch electrodes for measuring the resting membrane potential	65

4.5	Background currents which flow through voltage-gated channels open at resting membrane potential also participate in V_{rest}	67
	Further reading	67

Chapter 5 *The Voltage-Gated Channels of Na⁺ Action Potentials*

C. Hammond

5.1	Properties of action potentials	69
5.1.1	The different types of action potentials	69
5.1.2	Na ⁺ and K ⁺ ions participate in the action potential of axons	69
5.1.3	Na ⁺ -dependent action potentials are all or none and propagate along the axon with the same amplitude	71
5.1.4	Questions about the Na ⁺ -dependent action potential	71
5.2	The depolarization phase of Na ⁺ -dependent action potentials results from the transient entry of Na ⁺ ions through voltage-gated Na ⁺ channels	71
5.2.1	The Na ⁺ channel consists of a principal large α -subunit with four internal homologous repeats and auxiliary β -subunits	71
5.2.2	Membrane depolarization favours conformational change of the Na ⁺ channel towards the open state; the Na ⁺ channel then quickly inactivates	74
5.2.3	The time during which the Na ⁺ channel stays open varies around an average value, τ_o , called the mean open time	75
5.2.4	The $i_{\text{Na}}-V$ relation is linear: the Na ⁺ channel has a constant unitary conductance γ_{Na}	77
5.2.5	The probability of the Na ⁺ channel being in the open state increases with depolarization to a maximal level	77
5.2.6	The macroscopic Na ⁺ current (I_{Na}) has a steep voltage dependence of activation and inactivates within a few milliseconds	80
5.2.7	Segment S4, the region between segments S5 and S6, and the region between domains III and IV play a significant role in activation, ion permeation and inactivation, respectively	83
5.2.8	Conclusion: the consequence of the opening of a population of N Na ⁺ channels is a transient entry of Na ⁺ ions which depolarizes the membrane above 0 mV	87
5.3	The repolarization phase of the sodium-dependent action potential results from Na ⁺ channel inactivation and partly from K ⁺ channel activation	87
5.3.1	The K ⁺ channel consists of an α -subunit with a single repeat and auxiliary β -subunits	87
5.3.2	Membrane depolarization favours the conformational change of the delayed rectifier channel towards the open state	88
5.3.3	The open probability of the delayed rectifier channel is stable during a depolarization in the range of seconds	89
5.3.4	The K ⁺ channel has a constant unitary conductance γ_{K}	91
5.3.5	The macroscopic delayed rectifier K ⁺ current (I_{K}) has a delayed voltage dependence of activation and inactivates within tens of seconds	92
5.3.6	Conclusion: during an action potential the consequence of the delayed opening of K ⁺ channels is an exit of K ⁺ ions which repolarizes the membrane to resting potential	93
5.4	Sodium-dependent action potentials are initiated at the axon initial segment in response to a membrane depolarization and then actively propagate along the axon	94
5.4.1	Summary of the Na ⁺ -dependent action potential	95

5.4.2	Depolarization of the membrane to the threshold for voltage-gated Na^+ channel activation has two origins	95
5.4.3	The site of initiation of Na^+ -dependent action potentials is the axon initial segment	96
5.4.4	The Na^+ -dependent action potential actively propagates along the axon to axon terminals	97
5.4.5	Do the Na^+ and K^+ concentrations change in the extracellular or intracellular media during firing?	97
5.4.6	The role of the Na^+ -dependent action potential is to evoke neurotransmitter release	99
5.4.7	Characteristics of the Na^+ -dependent action potential are explained by the properties of the voltage-gated Na^+ channel	99
Appendix 5.1	Current clamp recording	99
Appendix 5.2	Voltage clamp recording	101
Appendix 5.3	Patch clamp recording	102
A5.3.1	The various patch clamp recording configurations	103
A5.3.2	Principles of the patch clamp recording technique	105
A5.3.3	The unitary current i is a rectangular step of current	107
A5.3.4	Determination of the conductance of a channel	107
A5.3.5	Mean open time of a channel	108
	Further reading	109

Chapter 6 *The Voltage-Gated Channels of Ca^{2+} Action Potentials: Generalization*

C. Hammond

6.1	Properties of Ca^{2+}-dependent action potentials	111
6.1.1	Ca^{2+} and K^+ ions participate in the action potential of endocrine cells	111
6.1.2	Questions about the Ca^{2+} -dependent action potential	111
6.2	The depolarizing or plateau phase of Ca^{2+}-dependent action potentials results from the transient entry of Ca^{2+} ions through voltage-gated Ca^{2+} channels	112
6.2.1	The voltage-gated Ca^{2+} channels are a diverse group of multisubunit proteins	113
6.2.2	The L, N and P-type Ca^{2+} channels open at membrane potentials positive to -20 mV; they are high-threshold Ca^{2+} channels	115
6.2.3	Macroscopic L, N and P-type Ca^{2+} currents activate at a high threshold and inactivate with different time courses	118
6.3	The repolarization phase of Ca^{2+}-dependent action potentials results from the activation of K^+ currents I_K and $I_{K(\text{Ca})}$	123
6.3.1	The Ca^{2+} -activated K^+ currents are classified as big K (BK) channels and small K (SK) channels	124
6.3.2	Ca^{2+} entering during the depolarization or the plateau phase of Ca^{2+} -dependent action potentials activates $\text{K}_{(\text{Ca})}$ channels	126
6.4	Calcium-dependent action potentials are initiated in axon terminals or in dendrites	128
6.4.1	Depolarization of the membrane to the threshold for the activation of L-, N- and P-type Ca^+ channels has two origins	128
6.4.2	The role of the calcium-dependent action potentials is to provide a local and transient increase of $[\text{Ca}^{2+}]_i$ to trigger secretion, contraction and other Ca^{2+} -gated processes	130
6.5	A note on voltage-gated channels and action potentials	131

Appendix 6.1	Fluorescence measurements of intracellular Ca^{2+} concentration	131
Y. Tan	A6.1.1 The interaction of light with matter	131
	A6.1.2 The return from the excited state	133
	A6.1.3 Fluorescence measurements: general points	134
	A6.1.4 Fluorescence imaging hardware	135
	A6.1.5 Methods of calcium measurement by fluorescence	135
	A6.1.6 Two-photon absorption	137
	A6.1.7 Measurement of other ions by fluorescence techniques	138
Appendix 6.2	Tail currents	139
Further reading		140

Chapter 7 *The Chemical Synapses*

C. Hammond

7.1	The synaptic complex's three components: presynaptic element, synaptic cleft and postsynaptic element	142
7.1.1	The pre- and postsynaptic elements are morphologically and functionally specialized	142
7.1.2	General functional model of the synaptic complex	144
7.1.3	Complementarity between the neurotransmitter stored and released by the presynaptic element and the nature of receptors in the postsynaptic membrane	146
7.2	The interneuronal synapses	149
7.2.1	In the CNS the most common synapses are those where an axon terminal is the presynaptic element	149
7.2.2	At low magnification, the axo-dendritic synaptic contacts display features implying various functions	149
7.2.3	Interneuronal synapses display ultrastructural characteristics that vary between two extremes: types 1 and 2.	150
7.3	The neuromuscular junction is the group of synaptic contacts between the terminal arborization of a motor axon and a striated muscle fibre	152
7.3.1	In the axon terminals, the synaptic vesicles are concentrated at the level of the electron-dense bars; they contain acetylcholine	153
7.3.2	The synaptic cleft is narrow and occupied by a basal lamina which contains acetylcholinesterase	153
7.3.3	Nicotinic receptors for acetylcholine are abundant in the crests of the folds in the postsynaptic membrane	155
7.3.4	Mechanisms involved in the accumulation of postsynaptic receptors in the folds of the postsynaptic muscular membrane	155
7.4	The synapse between the vegetative postganglionic neuron and the smooth muscle cell	156
7.4.1	The presynaptic element is a varicosity of the postganglionic axon	156
7.4.2	The width of the synaptic cleft is very variable	158
7.4.3	The autonomous postganglionic synapse is specialized to ensure a widespread effect of the neurotransmitter	158
7.5	Example of a neuroglandular synapse	159
7.6	Summary	160

Appendix 7.1	Neurotransmitters, agonists and antagonists	160
A7.1.1	Criteria to be satisfied before a molecule can be identified as a neurotransmitter	161
A7.1.2	Types of neurotransmitter	161
A7.1.3	Agonists and antagonists of a receptor	162
Appendix 7.2	Identification and localization of neurotransmitters and their receptors	162
M. Esclapez		
A7.2.1	Immunocytochemistry	162
A7.2.2	<i>In situ</i> hybridization	166
Further reading		167

Chapter 8 *Neurotransmitter Release*

C. Hammond

8.1	Observations and questions	169
8.1.1	Quantitative data on synapse morphology and synaptic transmission	169
8.1.2	Ways of estimating neurotransmitter release in central mammalian synapses	171
8.1.3	Questions	174
8.2	Presynaptic processes I: From presynaptic spike to $[Ca^{2+}]_i$ increase	174
8.2.1	The presynaptic Na^+ -dependent spike depolarizes the presynaptic membrane, opens presynaptic Ca^{2+} channels and triggers Ca^{2+} entry	174
8.2.2	Ca^{2+} enters the presynaptic bouton during the time course of the presynaptic spike through high-voltage-activated Ca^{2+} channels (N- and P/Q-types)	174
8.2.3	Presynaptic $[Ca^{2+}]_i$ increase is transient and restricted to micro- or nanodomains close to docked vesicles	175
8.2.4	Ca^{2+} clearance makes presynaptic $[Ca^{2+}]_i$ increase transient: it shapes its amplitude and duration	179
8.3	Presynaptic processes II: From $[Ca^{2+}]_i$ increase to synaptic vesicle fusion	180
8.3.1	Overview of the hypothetical vesicle cycle in presynaptic terminals	180
8.3.2	Docking: a subpopulation of synaptic vesicles is docked to the active zone close to Ca^{2+} channels by means of specific pairing of vesicular and plasma membrane proteins	182
8.3.3	Three to four Ca^{2+} ions must bind to Ca^{2+} receptor(s) to initiate vesicle fusion (exocytosis)	185
8.3.4	Fusion: from Ca^{2+} binding to exocytosis	186
8.3.5	Pharmacology of neurotransmitter release	187
8.4	Processes in the synaptic cleft: from transmitter release in the cleft to transmitter clearance from the cleft	188
8.4.1	The amount of neurotransmitter released in the synaptic cleft	188
8.4.2	Transmitter time course in the synaptic cleft is brief and depends mainly on transmitter binding to target proteins	190
8.5	Summary	191
Appendix 8.1	Quantal nature of neurotransmitter release	193
A8.1.1	Spontaneous release of acetylcholine at the neuromuscular junction evokes miniature endplate potentials: the notion of quanta	193
A8.1.2	The quantal composition of EPSPs and IPSPs	194
Appendix 8.2	The probabilistic nature of neurotransmitter release	194

A8.2.1	The neuromuscular junction as a model	194
A8.2.2	Inhibitory synapses between interneurons and the Mauthner cell in the teleost fish bulb, as a model	196
Further reading		198
 PART 2 Ionotropic and Metabotropic Receptors in Synaptic Transmission and Sensory Transduction		
 Chapter 9 <i>The Ionotropic Nicotinic Acetylcholine Receptors</i>		
C. Hammond		
9.1	Observations	
9.2	The torpedo or muscle nicotinic receptor of acetylcholine is a heterologous pentamer $\alpha_2\beta\gamma\delta$	202
9.2.1	Nicotinic receptors have a rosette shape with an aqueous pore in the centre	203
9.2.2	The four subunits of the nicotinic receptor are assembled as a pentamer $\alpha_2\beta\gamma\delta$	203
9.2.3	Each subunit presents two main hydrophilic domains and four hydrophobic domains	205
9.2.4	Each α -subunit contains one acetylcholine receptor site located in the hydrophilic NH_2 terminal domain	206
9.2.5	The pore of the ion channel is lined by the M2 transmembrane segments of each of the five subunits	207
9.3	Binding of two acetylcholine molecules favours conformational change of the protein towards the open state of the cationic channel	207
9.3.1	Demonstration of the binding of two acetylcholine molecules	207
9.3.2	The nicotinic channel has a selective permeability to cations: its unitary conductance is constant	208
9.3.3	The time during which the channel stays open varies around an average value τ_o , the mean open time, and is a characteristic of each nicotinic receptor	212
9.4	The nicotinic receptor desensitizes	214
9.5	nAChR-mediated synaptic transmission at the neuromuscular junction	217
9.5.1	Miniature and endplate synaptic currents are recorded at the neuromuscular junction	217
9.5.2	Synaptic currents are the sum of unitary currents appearing with variable delays and durations	219
9.6	Nicotinic transmission pharmacology	220
9.6.1	Nicotinic agonists	220
9.6.2	Competitive nicotinic antagonists	220
9.6.3	Channel blockers	221
9.6.4	Acetylcholinesterase inhibitors	222
9.7	Summary	223
Appendix 9.1	The neuronal nicotinic receptors	223
Further reading		225

Chapter 10 *The Ionotropic GABA_A Receptor*

C. Hammond

10.1	Observations and questions	227
10.2	GABA _A receptors are hetero-oligomeric proteins with a structural heterogeneity	227
10.2.1	The diversity of GABA _A receptor subunits	227
10.2.2	Subunit composition of native GABA _A receptors and their binding characteristics	228
10.3	Binding of two GABA molecules leads to a conformational change of the GABA _A receptor into an open state; the GABA _A receptor desensitizes	230
10.3.1	GABA binding site	230
10.3.2	Evidence for the binding of two GABA molecules	230
10.3.3	The GABA _A channel is selectively permeable to Cl ⁻ ions	230
10.3.4	The single-channel conductance of GABA _A channels is constant in symmetrical Cl ⁻ solutions, but varies as a function of potential in asymmetrical solutions	233
10.3.5	Mean open time of the GABA _A channel	234
10.3.6	The GABA _A receptor desensitizes	235
10.4	Pharmacology of the GABA _A receptor	237
10.4.1	Bicuculline and picrotoxin reversibly decrease total GABA _A current; they are respectively competitive and non-competitive antagonists of the GABA _A receptor	237
10.4.2	Benzodiazepines, barbiturates and neurosteroids reversibly potentiate total GABA _A current; they are allosteric agonists at the GABA _A receptor	238
10.4.3	β-carbolines reversibly decrease total GABA _A current; they bind at the benzodiazepine site and are inverse agonists of the GABA _A receptor	242
10.5	GABA _A -mediated synaptic transmission	243
10.5.1	The GABAergic synapse	243
10.5.2	The synaptic GABA _A -mediated current is the sum of unitary currents appearing with variable delays and durations	244
10.5.3	The consequences of the synaptic activation of GABA _A receptors depend on the relative values of E_{Cl} and V_m	246
10.5.4	What shapes the decay phase of GABA _A -mediated currents?	247
10.6	Summary	249
Appendix 10.1	Mean open time and mean burst duration of the GABA _A single-channel current	249
	Further reading	250

Chapter 11 *The Ionotropic Glutamate Receptors*

C. Hammond

11.1	The three different types of ionotropic glutamate receptors have a common structure and participate in fast glutamatergic synaptic transmission	251
11.1.1	Ionotropic glutamate receptors have the name of their selective or preferential agonist	251
11.1.2	The three ionotropic receptors participate in fast glutamatergic synaptic transmission	251
11.2	AMPA receptors are an ensemble of cationic receptor-channels with different permeabilities to Ca ²⁺ ions	253

11.2.1	The diversity of AMPA receptors results from subunit combination, alternative splicing and post-transcriptional nuclear editing	253
11.2.2	The native AMPA receptor is permeable to cations and has a unitary conductance of 8 pS	254
11.2.3	AMPA receptors are permeable to Na ⁺ , K ⁺ and Ca ²⁺ ions unless the edited form of GluR2 is present; in the latter case, AMPA receptors are impermeable to Ca ²⁺ ions	257
11.2.4	The presence of flip or flop isoforms plays a role on the amplitude of the total AMPA current	257
11.3	Kainate receptors are an ensemble of cationic receptor channels with different permeabilities to Ca²⁺ ions	257
11.3.1	The diversity of kainate receptors	257
11.3.2	Native kainate receptors are permeable to cations	258
11.4	NMDA receptors are cationic receptor-channels highly permeable to Ca²⁺ ions; they are blocked by Mg²⁺ ions at voltages close to the resting potential, which confers strong voltage-dependence	259
11.4.1	Molecular biology of NMDA receptors	259
11.4.2	Native NMDA receptors have a high unitary conductance of 40–50 pS	261
11.4.3	The NMDA channel is highly permeable to monovalent cations and to Ca ²⁺	261
11.4.4	NMDA channels are blocked by physiological concentrations of extracellular Mg ²⁺ ions; this block is voltage-dependent	262
11.4.5	Glycine is a co-agonist of NMDA receptors	265
11.4.6	Conclusions on NMDA receptors	265
11.5	Synaptic responses to glutamate are mediated by NMDA and non-NMDA receptors	267
11.5.1	Glutamate receptors are co-localized in the postsynaptic membrane of glutamatergic synapses	267
11.5.2	The glutamatergic postsynaptic current is inward and can have at least two components in the absence of extracellular Mg ²⁺ ions	268
11.5.3	The glutamatergic postsynaptic depolarization (EPSP) has at least two components in the absence of extracellular Mg ²⁺ ions	270
11.5.4	Synaptic depolarization recorded in physiological conditions: factors controlling NMDA receptors activation	270
11.6	Summary	272
	Further reading	272

Chapter 12 *Ionotropic Mechanoreceptors: the Mechanosensitive Channels*

C. Bourque

12.1	Mechanoreception in sensory neurons is associated with the production of a receptor potential	274
12.2	Discovery of mechanosensitive ion channels provided a potential molecular mechanism for mechanotransduction	274
12.3	Structural basis for the mechanical gating of ion channels	275
12.3.1	Intrinsic and extrinsic forms of mechanical gating	275
12.3.2	Channels regulated by coupling molecules oriented orthogonally to the membrane	275

12.3.3	Channels regulated by coupling molecules parallel to the membrane	275
12.3.4	Other gating configurations	276
12.4	Classification of stretch-sensitive ion channels	276
12.4.1	Patch clamp experiments reveal the existence of stretch-activated and stretch-inactivated channels	276
12.4.2	Ionic permeability of stretch-sensitive channels	277
12.5	Mechanosensitive ion channels and mechanotransduction	278
12.6	Osmoreceptors in the central nervous system	278
12.6.1	Electrical activity and neurohypophyseal hormone secretion	279
12.6.2	Magnocellular neurosecretory cells in the hypothalamus are intrinsic osmoreceptors	280
12.7	Osmoreception in magnocellular neurosecretory cells	281
12.7.1	Osmoreceptor potentials reflect the modulation of a non-selective cationic conductance	281
12.7.2	Changes in cell volume directly regulate the macroscopic cationic conductance in magnocellular neurosecretory cells	281
12.7.3	Magnocellular neurosecretory cells express stretch-inactivated cationic channels	282
12.7.4	The inhibitory effects of Gd^{3+} provide pharmacological evidence for the involvement of the stretch-inactivated cation channels in osmoreception	283
12.7.5	Molecular basis for mechanotransduction in osmoreceptors	283
12.8	Conclusions	284
	Further reading	286

Chapter 13 *The Metabotropic GABA_B Receptors*

D. Mott

13.1	GABA_B receptors were originally discovered because of their insensitivity to bicuculline and their sensitivity to baclofen	287
13.2	Structure of the GABA_B receptor	288
13.2.1	GABA _B receptors belong to family-3 G-protein coupled receptors	288
13.2.2	GABA _B receptors are heterodimers	291
13.2.3	GABA _B receptors are located throughout the brain at both presynaptic and postsynaptic sites	292
13.2.4	Summary	292
13.3	GABA_B receptors are G-protein-coupled to a variety of different effector mechanisms	292
13.3.1	GABA _B receptors are coupled to inhibitory G proteins	293
13.3.2	GABA _B receptors regulate the activity of adenylyl cyclase	294
13.3.3	GABA _B receptor activation inhibits voltage-dependent calcium channels	296
13.3.4	GABA _B receptors activate potassium channels	302
13.3.5	Summary	308
13.4	The functional role of GABA_B receptors in synaptic activity	308
13.4.1	Postsynaptic GABA _B receptors produce an inhibitory postsynaptic current	309
13.4.2	Presynaptic GABA _B receptors inhibit the release of many different transmitters	310
13.5	Summary	312
	Further reading	313

Chapter 14 *The Metabotropic Glutamate Receptors*

G. Bhavé and R. Gereau

14.1	What is the receptor underlying glutamate-stimulated PI hydrolysis? – The cloning of metabotropic glutamate receptor genes	314
14.2	How do metabotropic glutamate receptors carry out their function? – Structure–function studies of metabotropic glutamate receptors	315
14.3	What biochemical means do metabotropic glutamate receptors utilize to elicit physiological changes in the nervous system? – Signal transduction studies of metabotropic glutamate receptors	317
14.4	What are the functions of metabotropic glutamate receptors in the nervous system? – Physiological and genetic studies of mGluRs	319
14.5	How are metabotropic glutamate receptors specifically localized in neurons to execute their functions? – Studies of mGluR postsynaptic localization	323
14.6	How is the activity of metabotropic glutamate receptors modulated? – Studies of mGluR desensitization	324
14.7	Summary	325
	Further reading	325

Chapter 15 *The Metabotropic Olfactory Receptors*

C. Hammond

15.1	The olfactory receptor cells are sensory neurons located in the olfactory neuroepithelium	327
15.1.1	The olfactory neurons are bipolar cells that project to the olfactory bulb	327
15.1.2	Odorants diffuse through the extracellular mucous matrix before interacting with the chemosensory membrane of olfactory receptor neurons	329
15.2	The response of olfactory receptor neurons to odours is a membrane depolarization which elicits action potential generation	329
15.3	Odorants bind to a family of G-protein-linked receptors which activate adenylate cyclase	330
15.3.1	Odorant receptors are a family of G-protein-linked receptors	330
15.3.2	The activation of odorant receptors leads to the activation of adenylate cyclase and the rapid formation of cAMP via the activation of a G_s -like protein	331
15.4	cAMP opens a cyclic nucleotide-gated channel and generates an inward current	332
15.4.1	The olfactory cyclic nucleotide-gated channel is a ligand-gated channel composed of at least two different subunits	332
15.4.2	cAMP directly opens a cyclic nucleotide-gated channel	333
15.4.3	The activation of N cyclic nucleotide-gated channels evokes an inward depolarizing current carried by cations	338
15.4.4	The cyclic nucleotide-gated conductance and the odour-gated conductance are identical	342
15.4.5	Conclusions	342
15.5	The odorant-evoked inward current evokes a membrane depolarization that spreads electronically to the axon hillock where it can elicit action potentials	343

15.5.1	The odorant-induced inward current depolarizes the membrane of olfactory receptors: the generator potential	343
15.5.2	The odorant-induced depolarization takes place in the cilia; spikes are initiated in the soma-initial axon segment, and the pattern of spike discharge propagated codes for the concentration and duration of the odorant stimulus	344
15.5.3	The nature of the odorant would be coded by the nature of the olfactory neuron stimulated and the synaptic arrangements in the olfactory bulb	344
15.6	Conclusions	345
	Further reading	345

PART 3 Somato-Dendritic Processing and Plasticity of Postsynaptic Potentials

Chapter 16 *Somato-Dendritic Processing of Postsynaptic Potentials.*

I: Passive Properties of Dendrites

C. Hammond

16.1	Propagation of excitatory and inhibitory postsynaptic potentials through the dendritic arborization	350
16.1.1	The complexity of synaptic organization (Figure 16.1)	350
16.1.2	Passive decremental propagation of postsynaptic potentials	351
16.1.3	Passive and non-decremental propagation of postsynaptic potentials	351
16.2	Summation of excitatory and inhibitory postsynaptic potentials	351
16.2.1	Linear and nonlinear summation of excitatory postsynaptic potentials	351
16.2.2	Linear and nonlinear summation of inhibitory postsynaptic potentials	354
16.2.3	The integration of excitatory and inhibitory postsynaptic potentials partly determines the configuration of the postsynaptic discharge	354
16.3	Summary	354
	Further reading	355

Chapter 17 *Subliminal Voltage-Gated Currents of the Somato-Dendritic Membrane*

C. Hammond

17.1	Observations and questions	359
17.2	The subliminal voltage-gated currents that depolarize the membrane	359
17.2.1	The persistent inward Na^+ current, I_{NaP}	359
17.2.2	The low-threshold transient Ca^{2+} current, I_{CaT}	361
17.2.3	The hyperpolarization-activated cationic current, I_h , I_p , I_Q	364
17.3	The subliminal voltage-gated currents that hyperpolarize the membrane	367
17.3.1	The rapidly inactivating transient K^+ current: I_A or I_{Af}	367
17.3.2	The slowly inactivating transient K^+ current, I_D or I_{As}	368
17.3.3	The K^+ currents activated by intracellular Ca^{2+} ions, I_{KCa}	369
17.3.4	The K^+ current sensitive to muscarine, I_M	369

17.3.5	The inward rectifier K^+ current, I_{Kir}	371
17.4	Conclusions	372
	Further reading	372

Chapter 18 *Somato-Dendritic Processing of Postsynaptic Potentials.*

II. Role of Subliminal Depolarizing Voltage-Gated Currents

C. Hammond

18.1	Persistent Na^+ channels are present in soma and dendrites of neocortical neurons; I_{NaP} boosts EPSPs in amplitude and duration	375
18.1.1	Persistent Na^+ channels are present in the dendrites and soma of pyramidal neurons of the neocortex	375
18.1.2	Dendritic persistent Na^+ channels are activated by EPSPs; in turn, I_{NaP} boosts EPSP amplitude	376
18.2	T-type Ca^{2+} channels are present in dendrites of neocortical neurons; I_{CaT} boosts EPSPs in amplitude and duration	378
18.2.1	T-type Ca^{2+} channels are present in dendrites of pyramidal neurons of the hippocampus	378
18.2.2	Dendritic T-type Ca^{2+} channels are activated by EPSPs; in turn, I_{CaT} boosts EPSPs amplitude	378
18.3	The hyperpolarization-activated cationic current I_h is present in dendrites of hippocampal pyramidal neurons; for EPSPs, dendritic I_h decreases the current transmitted from the dendrites to the soma	382
18.3.1	H-type cationic channels are expressed in dendrites of pyramidal neurons of the hippocampus	382
18.3.2	Dendritic H-type cationic channels are activated by IPSPs; in turn, I_h decreases EPSPs amplitude	383
18.4	Functional consequences	386
18.4.1	Amplification of distal EPSPs by I_{NaP} and I_{CaT} counteracts their attenuation owing to passive propagation to the soma; it also favours temporal summation versus spatial summation	386
18.4.2	Activation of dendritic I_{CaT} generates a local dendritic $[Ca^{2+}]_i$ transient	386
18.4.3	Activation of dendritic I_h , I_{NaP} and I_{CaT} alter the local membrane resistance and time constant	386
18.5	Conclusions	386
	Further reading	387

Chapter 19 *Somato-Dendritic Processing of Postsynaptic Potentials.*

III. Role of High-Voltage-Activated Depolarizing Currents

C. Hammond

19.1	High-voltage-activated Na^+ and/or Ca^{2+} channels are present in the dendritic membrane of some CNS neurons, but are they distributed with comparable densities in soma and dendrites?	390
19.1.1	High-voltage-activated Na^+ channels are present in some dendrites	390

19.1.2	Dendritic Na ⁺ channels are opened by EPSPs and the resultant Na ⁺ current boosts EPSPs in amplitude and duration	392
19.1.3	Dendritic Na ⁺ channels are opened by backpropagating Na ⁺ action potentials	394
19.2	High-voltage-activated Ca²⁺ channels are present in the dendritic membrane of some CNS neurons, but are they distributed with comparable densities in soma and dendrites?	398
19.2.1	High-voltage-activated Ca ²⁺ channels are present in some dendrites	398
19.2.2	High-voltage-activated Ca ²⁺ channels of Purkinje cell dendrites are opened by climbing fibre EPSP; this initiates Ca ²⁺ action potentials in the dendritic tree of Purkinje cells	400
19.2.3	Dendritic high-voltage-activated Ca ²⁺ channels are opened by backpropagating Na ⁺ action potentials	403
19.3	Functional consequences	403
19.3.1	Amplification of distal synaptic responses by dendritic HVA currents counteracts their attenuation due to passive propagation to the soma	403
19.3.2	Active backpropagation of Na ⁺ spikes in the dendritic tree depolarizes the dendritic membrane, with multiple consequences	404
19.3.3	Initiation of Ca ²⁺ spikes in the dendritic tree of Purkinje cells evokes a widespread intradendritic [Ca ²⁺] increase	405
19.4	Conclusions	405
	Further reading	406

Chapter 20 *Firing Patterns of Neurons*

C. Hammond

20.1	Medium spiny neurons of the neostriatum are silent neurons that respond with a long latency	407
20.1.1	Medium spiny neurons are silent at rest owing to the activation of an inward rectifier K ⁺ current	407
20.1.2	When activated, the response of medium spiny neurons is a long-latency regular discharge	409
20.2	Inferior olivary cells are silent neurons that can oscillate	410
20.2.1	Inferior olivary cells are silent at rest in the absence of afferent activity	410
20.2.2	When depolarized, inferior olivary cells oscillate at a low frequency (3–6 Hz)	411
20.2.3	When hyperpolarized, inferior olivary cells oscillate at a higher frequency (9–12 Hz)	413
20.3	Purkinje cells are pacemaker neurons that respond by a complex spike followed by a period of silence	414
20.3.1	Purkinje cells present an intrinsic tonic firing that depends on a persistent Na ⁺ current	414
20.3.2	Purkinje cells respond to climbing fibre activation by a complex spike	414
20.4	Thalamic and subthalamic neurons are pacemaker neurons with two intrinsic firing modes: a tonic and a bursting mode	417
20.4.1	The intrinsic tonic (single-spike) mode depends on a persistent Na ⁺ current	419
20.4.2	The bursting mode depends on a cascade of subliminal inward currents: I_h , I_{CaT} , I_{CaN}	420
20.4.3	The transition from one mode to the other in response to synaptic inputs	420
	Further reading	423

Chapter 21 *Synaptic Plasticity*

C. Hammond

21.1	Short-term potentiation (STP) of a cholinergic synaptic response as an example of short-term plasticity: the cholinergic response of muscle cells to motoneuron stimulation	424
21.2	Long-term potentiation (LTP) of a glutamatergic synaptic response: example of the glutamatergic synaptic response of pyramidal neurons of the CA1 region of the hippocampus to Schaffer collaterals activation	425
21.2.1	The Schaffer collaterals are axon collaterals of CA3 pyramidal neurons which form glutamatergic excitatory synapses with dendrites of CA1 pyramidal neurons	425
21.2.2	Observation of the long-term potentiation of the Schaffer collateral-mediated EPSP	427
21.2.3	Long-term potentiation (LTP) of the glutamatergic EPSP recorded in CA1 pyramidal neurons results from an increase of synaptic efficacy (or synaptic strength)	427
21.2.4	Induction of LTP results from a transient enhancement of glutamate release and a rise in postsynaptic intracellular Ca^{2+} concentration	429
21.2.5	Expression of LTP (also called maintenance) involves a persistent enhancement of the AMPA component of the EPSP	434
21.2.6	Multiple ways to induce LTP, multiple forms of LTP and multiple ways to block LTP induction	435
21.2.7	Summary: principal features of LTP in the Schaffer collateral–pyramidal cell glutamatergic transmission	437
21.3	The long-term depression (LTD) of a glutamatergic response: example of the response of Purkinje cells of the cerebellum to parallel fibre stimulation	437
21.3.1	The long-term depression of a postsynaptic response (EPSC or EPSP) is a decrease of synaptic efficacy	437
21.3.2	Induction of LTD requires a rise in postsynaptic intracellular Ca^{2+} concentration and the activation of postsynaptic AMPA receptors	438
21.3.3	The expression of LTD involves a persistent desensitization of postsynaptic AMPA receptors	443
21.3.4	Second messengers are required for LTD induction: examples of protein kinases C and nitric oxide (NO)	443
21.3.5	The different ways to induce or block cerebellar LTD	446
21.3.6	Summary: Principal features of LTD in parallel-fibre/Purkinje-cell glutamatergic transmission	448
	Further reading	448

PART 4 Activity and Development of Networks: The Hippocampus as an Example

Chapter 22 *The Adult Hippocampal Network*

C. Hammond

22.1	Observations and questions	451
22.2	The hippocampal circuitry	453
22.2.1	Ammon's horn	453
22.2.2	The dentate gyrus	454
22.2.3	Principal cells form a tri-neural excitatory circuit	457
22.2.4	Extrinsic afferences to principal cells and interneurons	457

22.3	Activation of interneurons evoke inhibitory GABAergic responses in postsynaptic pyramidal cells	458
22.3.1	Experimental protocol to study pairs of neurons	458
22.3.2	Unitary inhibitory postsynaptic currents (IPSCs) evoked by different types of interneurons are all GABA _A -mediated but have different kinetics when recorded at the level of the soma	458
22.3.3	GABA _A -mediated IPSCs generate IPSPs in postsynaptic pyramidal cells	461
22.3.4	GABA _B -mediated IPSPs are also recorded in pyramidal neurons in response to strong interneuron stimulation	464
22.4	Activation of principal cells evokes excitatory glutamatergic responses in postsynaptic interneurons and other principal cells (synchronization in CA3)	464
22.4.1	Pyramidal neurons evoke AMPA-mediated EPSPs in interneurons	465
22.4.2	EPSPs in interneurons lead to feedback inhibition of pyramidal neurons	466
22.4.3	CA3 pyramidal neurons are monosynaptically connected via glutamatergic synapses	466
22.4.4	Overview of intrinsic hippocampal circuits	466
22.5	Oscillations in the hippocampal network: example of sharp waves (SPW)	466
22.6	Summary	470
	Further reading	471

Chapter 23 *Maturation of the Hippocampal Network*

Y. Ben Ari and C. Hammond

23.1	GABAergic neurons and GABAergic synapses develop prior to glutamatergic ones	472
23.1.1	GABAergic interneurons divide and arborize prior to pyramidal neurons and granular cells	472
23.1.2	GABAergic synapses are the first synapses established on to pyramidal cells	473
23.1.3	Sequential expression of GABA and glutamate synapses is also observed in the hippocampus of subhuman primates <i>in utero</i>	474
23.1.4	Questions about the sequential maturation of GABA and glutamate synapses	476
23.2	GABA _A - and GABA _B -mediated responses differ in developing and mature brains	476
23.2.1	Activation of GABA _A receptors is depolarizing and excitatory in immature networks because of a high intracellular concentration of chloride	476
23.2.2	GABA _B -receptor-mediated IPSCs have a delayed expression in immature neurons	479
23.3	Network-driven giant depolarizing potentials (GDPs) provide most of the synaptic activity in the neonatal hippocampus	480
23.3.1	Giant depolarizing potentials result from GABAergic and glutamatergic synaptic activity	481
23.3.2	Giant depolarizing potentials are generated in the septal pole of the immature hippocampus and then propagate to the entire structure	482
23.4	Hypotheses on the role of the sequential expression of GABA- and glutamate-mediated currents and of giant depolarizing potentials	483
23.5	Conclusions	483
	Further reading	484
	Index	485